Dosing Guidelines for Common Medications

Products come in different formulations and doses. Read packages carefully to determine the specific concentration of liquids and/or the specific dose of each chewable or tablet. Always use the measuring device packaged with the medication; do not substitute another measuring device.

Fever reducing medications: Please DO NOT alternate ibuprofen and acetominophen ~ this can easily lead to overdose!

Ibuprofen (Motrin, Advil)

Indication:

- Give for fever or pain
- Not recommended for relief of cold symptoms
- Not recommended for children under the age of 6m

Timing:

- You may give a dose of ibuprofen every 6-8 hours as needed
- Maximum doses in 24 hours: 4

Formulas:

Concentrated Infant drops 50mg/1.25ml
Children's liquid 100mg/5ml
Junior Strength Chewables 100mg each

• Adult Tablet/Capsules 200mg each

Weight	Drops	Liquid	Chewable	Adult tablet
12-17 lbs	1.25 ml	$\frac{1}{2}$ tsp = 2.5ml		
18-23 lbs	1.875 ml	¾ tsp = 3.75ml		
24-35 lbs	2.5ml (1.25ml x2)	1 tsp = 5 ml	1 chewable	
36-47 lbs	3.75 ml (1.875 x 2)	1 ½ tsp = 7.5ml	1 ½ chewables	
48-59 lbs		2 tsp = 10 ml	2 chewables	1 adult tab
60-71 lbs		2 ½ tsp = 12.5ml	2 ½ chewables	1 ½ adult tablet
72-95 lbs		3 tsp = 15ml	3 chewables	1 ½-2 tablets

Acetominophen (Tylenol)

Indication:

- Fever or pain
- Not recommended for relief of cold symptoms
- Not recommended under age 2m unless doctor advised

Timing:

- You may give acetominophen every 4-6 hours as needed
- Maximum number of doses in 24 hours: 5

Formulas:

• Children's liquid 160mg/5ml = 160mg/1 tsp

• Children's Chewables 80mg each

• Junior Chewables 160mg each

Weight	Liquid	80mg Chews	160mg Chews
6-11 lbs	½ tsp = 1.25ml		
12-17 lbs	½ tsp = 2.5 ml		
18-23 lbs	$\frac{34}{4} \text{ tsp} = 3.75 \text{ ml}$		
24-35 lbs	1 tsp = 5ml	2 chewables	1 chewable
36-47 lbs	$1 \frac{1}{2} \text{ tsp} = 7.5 \text{ml}$	3 chewables	1 ½ chewable
48-59 lbs	2 tsp = 10ml	4 chewables	2 chewables
60-71 lbs	2 ½ tsp=12.5 ml	5 chewables	2 ½ chewables
72-95 lbs	3 tsp = 15 ml	6 chewables	3 chewables

Diphenhydramine (Benadryl)

Indication:

- Hives, allergic reactions, itching or seasonal allergies
- Not recommended as a sleep aid
- Not recommended under age 6m unless doctor advised

Timing:

• You may give diphenhydramine every 6 hours as needed

Formulas:

• Children's liquid 12.5 mg/5 ml = 12.5 mg/1 tsp

• Children's Chews 12.5 mg each

Weight	Children's liquid	Children's Chewable	
10-16 lbs	½ tsp = 2.5ml		
17-21 lbs	3⁄4 tsp = 3.75 ml		
22-32 lbs	1 tsp = 5 ml	1 chewable	
33-42 lbs	1 ½ tsp = 7.5 ml	1 ½ chewables	
43-53 lbs	2 tsp = 10 ml	2 chewables	
54-64 lbs	2 ½ tsp = 12.5ml	2 ½ chewables	
65-75 lbs	3 tsp = 15 ml	3 chewables	
76-86 lbs	3 ½ tsp = 17.5ml	3 ½ chewables	

Certrizine(Zyrtec)

Indication:

- Hives, itching or seasonal allergy symptoms
- Not recommended as a sleep aid or for relief of cold symptoms unless otherwise advised by doctor
- Not recommended under 6m unless doctor advised

Timing:

• You may give certrizine(Zyrtec) 1-2x daily

Formulas:

Children's liquid 5mg/5ml = 5mg/1 tsp
Children's Chewable 5mg OR 10mg each
Adult tablets 5mg OR 10mg each

Age	Liquid	5mg dose	10mg dose
6-11 months	½ tsp = 2.5ml		
12-23 months	$\frac{1}{2}$ tsp = 2.5ml 1-2 x daily		
2-5 years	1 tsp daily	1 tablet daily	
6-11 years	1-2 tsp daily	1-2 tablets daily	1 tablet daily
12+ years	2 tsp daily	2 tablets daily	1 tablet daily